

نماذج مزاولة مهنة بكلوريوس طب عام

جزء 1

نسخة محلولة

EXAM

تنسيق وإعداد :

By: ASEEL Ali YOUS

د. أصيل علي يوس

اضغط على الرابط لمتابعة قناتنا على التلجرام

<https://t.me/medical02017>

ت: +٩٦٧ ٧٣٣٤٤٣٨٤٦

حقوق الطبع محفوظة لدى:

مكتبة حيكان

للمستلزمات القرطاسية والإلكترونية والتصوير

صنعاء - شارع حدة - جولة ريماس

أمام بوابة المجلس الطبي - جوار الياباني للصرافة

ت: +٩٦٧ ٧٧٥٧٥٥٠٧٠



1.Prescription Must contain all of the following except:

- a. Name of the patient
- b. Diagnosis
- c. Single or Married*
- d. Dosage form

2.If the Prescription is not clear:

- a. Return it to the Doctor*
- b. Give it to the Pharmacist
- c. Dispense the clear drugs
- d. Return it to the patient

3.which of the following drugs not used in Acute pain?

- a. Morphine
- b. Phentanyl
- c. Naloxone*
- d. Dexamethasone

4.which of the following is parenteral route of administration :

- a. subcutaneous
- b. IM
- c. IV
- d. All of the above*

5. Insulin Storage at:

- a. 0c
- b. 4-8c*
- c. 24c
- d. 25c

6.Which of the following β blocker used in ophthalmic preparation?

- a. Timolol*
- b. Propranolol
- c. Labetalol
- d. Pindolol

7.Fluxetine belongs to:

- a. Proton pump inhibitor
- b. ACEI
- c. Selective Serotonin reuptake inhibitor*
- d. Oral hypoglycemic

8.Treatment of Hypertension in pregnant woman:

- a. Captopril
- b. Verapamil
- c. Methyldopa*
- d. All above

9.dose of Klacid (Clarithromycin):

- a. three times per day
- b. twice per day
- c. once
- d. four times per day (1-2g divided in 2-4 times [250-500mg/tab])

9.Atenolol sold in KSA under trade name of:

- a. Indral
- b. Tenormin*
- c. Capoten
- d. Zovirax

10.Atenolol is:

- a. alpha blocker
- b. non selective beta blocker
- c. selectivebeta1 blocker *
- d. beta agonist

11.trade name of Fosinopril is:

- a. Capoten
- b. Reinetic
- c. Zestril
- d. Monopril*

12.Heparin is:

- a. Anti plattlet
- b. Anti coagulant*
- c. Fibrinolytic
- d. Analgesic

13.Advantage of Salbutamol over other bronchodilator:

- a. Longacting
- b. Morepotent
- c. Low Side effect
- d. Fast Acting*

14.Which of the following drugs are not OTC:

- a. Aspirin
- b. Bisacodyl
- c. Tegretol (carbamazepin)*
- d. Malox (antacid)

15. Side effect of oral Hypoglycemia:

- a. Hyperglycemia
- b. Hypoglycemia*
- c. Hypernatremia
- d. Hypertension

16. Generic name of Zantac:

- a. Rantidine*
- b. Cimitidine
- c. Famotodine
- d. Nizatidine

17. Diltiazim is:

- a. beta blocker
- b. Ca channel blocker *
- c. Na channel blocker
- d. Alph blocker

18. Over dose of panadol can be treated with:

- a. Naloxone
- b. EDTA
- c. Vitamin K
- d. Acetyl Cystiene*

19. qod means:

- a. three times daily
- b. everyday
- c. four times daily
- d. every other day*

20. qid means:

- a. three times per day
- b. two times per day
- c. four times per day*
- d. once daily

21. bid means:

- a. three times per day
- b. two times per day*
- c. four times per day
- d. once daily

22. tid means:

- a. three times per day*
- b. two times per day
- c. four times per day
- d. once daily

23. about Tetracycline all of the following are right except:

- a. It can suppress the growth of bacteria
- b. It binds to bacterial Ribosome
- c. active against G+ve and G-ve
- d. safe for children use with out affecting color of teeth*

24. All of these drugs are controlled drugs except:

- a. morphine
- b. Nalbuphine
- c. Aspirin*
- d. Methadone

25. Opioid toxicity can be treated with:

- a. Paracetamol
- b. Naloxone*
- c. Deferroxamin
- d. Vitamin K

26. Dapson is:

- a. Beta lactam Antibiotic
- b. Cephalosporines
- c. Sulphadrug*
- d. Antacid

27. Rheumatoid can be treatment with:

- a. Gold derivatives drugs *
- b. Platinum derivatives drugs
- c. Silver derivatives drugs
- d. Non of the above

28. Concentration of normal saline :

- a. 9%
- b. 5.3%
- c. 0.9%*
- d. 90%

29. which of the following dosage form can be administrated orally:

- a. spray
- b. Suppositories
- c. Ointment
- d. Elixir*

30. Aspirin used as:

- a. Analgesic
- b. Antipyretic
- c. Antinflammatory
- d. All above *

31. which of the following drugs inhibit liver enzymes :

- a. Paracetamol
- b. Epanutine(phenytoin)
- c. Cimitidine*
- d. Aspirin

33-Which of the following drugs Induce liver enzymes:

- a. Rifamicin*
- b. Rantidine
- c. Cimitidine
- d. Acyclovair

34. Side effect of Atropine:

- a. Diarrhea
- b. Headache
- c. Drymouth*
- d. Constipation*

35.Amphotricine is:

- a. Antibiotic
- b. Antifungal*
- c. Antiviral
- d. Antiprotozoal

36.Vassoprossine used in treatment of:

- a. Diabetes mellitus
- b. Diabetes Insipidus*
- c. Hyperthyrodism
- d. Hypothyrodism

37.Maximum dose of Diclophenac ((Voltaren)):

- a. 1000 mg/day
- b. 200 mg/day*
- c. 4000 mg/day
- d. 3200 mg/day

38. Side effect of Buscopan:

- a. Diarrhea
- b. Headache
- c. Drymouth*
- d. Constipation*

39.Metphormine used as:

- a. Antibiotic
- b. Antihypertensive
- c. Oral hypoglycemic*
- d. Antianginal

40.pseudo membranous colitis treated with:

- a. omeprazol
- b. mebendazol
- c. metronidazol*
- d. Amoxil

41. IM administration Site of Injection:

- a. Arm
- b. Restus Lateral (lateral restus in the eye)
- c. Buttocks
- d. All of the above

42. the Latin abbreviation for four times :

- a. qid*
- b. bid
- c. tid
- d. qd

43. Zinnat is:

- a. First generation cephalosporin's
- b. Quinolon
- c. Aminoglycosides
- d. Second generation cephalosporin's*

44. which of the following is not beta Agonist:

- a. Salbutamol
- b. Salmeterol
- c. Pindolol*
- d. Albutrol(Salbutamol)

45. Amoxicillin + clavulanic present in:

- a. Ciprobay
- b. Augmentin*
- c. Zantac
- d. Kl acid

46. Warfarin over dose can be treated with:

- a. Vitamin A
- b. Acetylcustiene
- c. Protamin sulphate
- d. Vitamin K*

47. Quality control is control of:

- a. condition
- b. specification
- c. shelf life
- d. All above *

48. Which of the following effect is not considered information of suspension:

- a. Use of chelating agent*
- b. of whiting agent*
- c. particle size
- d. Non of the above

49. Streptokinase is indicated for:

- a. Pulmonary embolism*
- b. Impaired fat absorption
- c. Tuberculosis
- d. Neoplastic anemia

49- Analgesic indicated for Asthmatic patient:

- a. Paracetamol*
- b. Aspirin
- c. Brufen
- d. Salbutamol

50- Antibiotic safe for Pregnant woman:

- a. Ciprofloxacin
- b. Gentamycin
- c. Amoxicillin*
- d. Tetracycline

51- Atenolol belongs to:

- a. NSAID
- b. AB
- c. β blocker *
- d. ACEI

52- Normal Potassium level in blood:

- a. 50-30
- b. 150-130
- c. 3.5-5 (3.5-5.0(mEq/L)).*
- d. 5.5-8.5

53- content of patient file is:

- a. name+age+gender+Diagnosis +medication*
- b. name+Diagnosis medication only
- c. name+age+gender+telephone

54- used in treatment of AIDS:

- a. Acyclovir*
- b. Rifampicin
- c. Zidovudine
- d. Prednisolone

55- Diclofenac taken:

- a. Before meal
- b. After meal*
- c. With in meal
- d. Not affected by food content

56-Neostigmine should stopped before Pregnancy by: ?

- a. Two weeks*
- b. Two months
- c. Two Years
- d. non above

57-Which of the following OTC:

- a. Tegretol
- b. Celebrex (Celecoxib=NSAID)*
- c. Augmentine
- d. Phenobarbital
- e. non above

58-Which of the following are oral aminoglycosides

- a. Gentamycin
- b. Amikacin
- c. Topramycin
- d. Neomycin*

59-Similar to Teico planin in the effect on G+ve bacteria:

- a. Jamica
- b. Gentamycin
- c. Kanamycin
- d. Vancomycin*

60-Which of the following is not Tetracycline?

- a. Minocycline
- b. Demeclocline
- c. Dilitiazem*
- d. Doxeycline

61-Which of the following are not Ca channel blockers?

- a. Amlodipine
- b. Cinnarazine*
- c. Verapamil
- d. Diltiazem

62-Which of the following are used in treatment of Hypertension ?

- a. Folicacid
- b. Amoxil
- c. Inderal*
- d. Amantadin

63-Spironolactone is:

- a. ACEI
- b. Antibiotic
- c. Antiviral
- d. Diuretics*

64-Which of the following Anti hypertensive are safe during Pregnancy:

- a. Nifedipine
- b. Amlodipine
- c. Methyldopa*
- d. Capoten

65- used intreatment of Amoebia giardia:

- a. yomesan
- b. Flagyl*
- c. Prazaquantil
- d. Mgsulphate

66-Scientific name of vitamin B1:

- a. Riboflavin
- b. Tochopherol
- c. Thiamine*
- d. Pyridoxine

67-Which of the following is contraindicated in Asthmatic patient :

- a. Ranitidine
- b. Aspirin*
- c. Atrovent
- d. Salbutamol

68-Period (duration) of treatment of Tuberculosis (TB):

- a. from 6 months to 9 months*
- b. from 6 weak to 9weak
- c. from 1 year to 1year
- d. from 2 year to 2year

69-OTC means:

- a. Over the content
- b. Over the counter Drugs *
- c. On the counter
- d. Ophthalmic Drugs

70-Antidote for Warfarine:

- a. Deferroxamin
- b. Protamin
- c. Vitamin K*
- d. Heparin

71-Antidote for INH:

- a. Naloxone
- b. Protamin sulphate
- c. Vitamin B6*
- d. Vitamin K

72-Generic name of Capoten :

- a. Ramipril
- b. Captopril*
- c. Lisinopril
- d. Propranolol

73-Antidote for Heparin:

- a. Deferroxamin
- b. Protamin*
- c. Vitamin K
- d. Naloxone

74-Generic name of Prozac:

- a. Imepamine
- b. Fluxotine*
- c. Citalopram
- d. Sertraline

75-Content of ORS:

- a. NaCl + Nalactate + Kcitate
- b. Glucose + NaCl + Nacitrate + Kcitate
- c. NaCl + Nalactate + Kcitate + MgOH
- d. Glucose+fructose+NaCl+KCl+Nacitrate+Kcitate (Glucose+NaCl+KCl+Na. citrate or bicarbonate)*

76-Ideal Temperature for Insulin storage:

- a. 15-20C
- b. 2-8C*
- c. 0-5C
- d. 2-4C

78-Theophyllin is:

- a. Bronchodilator*
- b. β 2 agonist
- c. Anticholinergic
- d. All of the above

79-Normal Hemoglobin level in blood:

- a. in male 15-20 in woman 25-30
- b. in male 13.5-16.5 in woman 12.5-15.5*
- c. 12-15
- d. in male 10-12 in woman 8-10

80-Antidote for Iron:

- a. Deferroxamin*
- b. Protamin
- c. Vitamin K
- d. Naloxone

81-Malaria Transmitted via:

- a. pugs
- b. Mosquito*
- c. Bacteria
- d. Fungus

82-Acyclovir used intreatment of:

- a. bacterial infection
- b. Viral infection*
- c. Nesseria
- d. Fungal infection

83-Prescription contains Prednsolone as:

- a. 1 X Tid X 3days
- b. 1 X Bid X 3days
- c. 1 X Qd X 3days

84-How many tablets will you give to the patient: ?

- a. 9
- b. 36
- c. 27
- d. 18*

84-Scientific name of Vitamin C:

- a. Acetyl Salycilic acid
- b. Ascorbic Acid*
- c. Cyanocobalamine
- d. Nicotenic Acid

85- used intreatment of Ascaris:

- a. Clindamycin
- b. Prazaquantil
- c. Metronidazol
- d. Mebendazole*

86-Vaccines Storage:

- a. Awayfromlight
- b. atRometemp.
- c. atrefrigerator*
- d. All above

87-The way that you deliver 100% of the drug to the blood (highest Bioavailability) and rapidaction:

- a. IM
- b. IV*
- c. Rectally
- d. Inhalation

88-Maximum dose of Paracetamol:

- a. 4gm/day*
- b. 40mg/day
- c. 2400mg/day
- d. 3tab/day

89-Use of Bariumsulphate:

- a. for Acne
- b. for Constipation
- c. Radiocontrast agent*
- d. for Diarrhea

90-Salivation Enzyme converts food into:

- a. Glucose
- b. Maltose*
- c. Fructose
- d. Sucrose

91-Maximum dose of captopril:

- a. 125mg/day
- b. 150mg/day*
- c. Threetab/day(450mg/day)25-125mg bid or tid
- d. 4gm/day

92-To avoid more absorption of toxic drugs use:

- a. saline
- b. Charcoal*
- c. Antibiotic
- d. Antidote

93-Type II diabetes use:

- a. Insulin
- b. Sulphonyl urea*
- c. H2 antagonist
- d. Glucose

94-RedMan Syndrome is the Side effect of:

- a. Aspirin
- b. Metronidazole
- c. Vancomycin*
- d. Gentamycin

95-TypeI diabetes use:

- a. oral hypoglycemic
- b. Insulin *
- c. Dextrosesol
- d. sulphonylurea

96-Which is contraindicated during pregnancy?

- a. Mythildopa
- b. Paracetamol
- c. Ciprofloxacin*
- d. Penicillin

97-Nephrotoxicity is the Side effect of:

- a. Paracetamol
- b. Metronidazole
- c. Vancomycin
- d. Gentamycin*

98-Hepato toxicity is the Side effect of:

- a. Paracetamol*
- b. Metronidazole
- c. Vancomycin
- d. Gentamycin

99-Generic name of Voltaren:

- a. Diclofenac Sodium*
- b. Piroxicam
- c. Acetylsalicylic acid
- d. Indomethacin

100-Ondansteron HCl is used for:

- a. Hypertension
- b. angina pectoris
- c. prophylaxis of migraine
- d. anti-emetic*

101- All of the severe Side effect of Chloramphenicol except:

- a. irreversible aplastic anaemia
- b. opticneuritis
- c. peripheralneuritis
- d. iron deficiency anaemia*

102-Ticlopidine is used for:

- a. anticoagulant
- b. antibiotic
- c. anticholinergic
- d. antiplatelets*

103-Teicoplanin is used for:

- a. anticoagulant
- b. antibiotic*
- c. anticholinergic
- d. antiplatelets

104-Retrovir is the brand name of:

- a. Famciclovir
- b. Zidovudine*
- c. Acyclovir
- d. Amantadine

105-Pseudo membranous colitis is the Side effect of:

- a. Vancomycin
- b. Flagyl
- c. Dapsone.
- d. Clindamycin*

106-Sporanoxis used to treat:(itraconazole)

- a. antifungal*
- b. antiviral
- c. antiprotozoal
- d. antibiotic

107-Fasomaxis the trade name of:

- a. Dapsone
- b. Amantadine
- c. Alendronate* (bisphosphonate)
- d. Warfarin

108-Bisphosphonate is used to treat:

- a. Acne
- b. Osteoporosis*
- c. Contraceptive
- d. constipation

109-The antacid that cause diarrhea is:

- a. Ca
- b. Al
- c. Mg*
- d. All of the above

110-Sodium Stibogluconate is used to treat:

- a. Leshmaniasis*
- b. Amoebiasis
- c. Ascaris
- d. Taenia

111-Amenorrhoea is defined as:

- a. painful menstruation
- b. Premenstrual syndrome
- c. Impotence
- d. Absence of menstruation*

112-What is the major complication of morphine over dose?

- a. AV-Block
- b. Respiratory depression*
- c. Cardiac Arrest
- d. Liver dysfunction

113-Which of the following is IN CORRECT?

- a. A syrup is an aqueous solution that has a sweet taste.
- b. Sustained release tablets should be crushed before dissolving when given to a patient*.
- c. Elixir is a solution that contains alcohol as diluent.
- d. Regular tablet is a solid dosage form.

114-The duration for treatment of Brucellosis is:

- a. 2weeks
- b. 2months
- c. 6weeks*
- d. 6months

115-SNF is:

- a. Southern Non-Fatal Fever
- b. Sinusitis Nasal Formula
- c. Saudi National Formulary*
- d. Computer software contains information about drugs, diseases and other scientific materials

116-In case of acetaminophen poisoning, the recommended antidote is:

- a. IV normal saline
- b. N-acetylcysteine *
- c. Epinephrine
- d. Activated charcoal

117-Which one of these drugs is NOT an antiarrhythmic agent?

- a. Procainamide
- b. Amitriptyline*
- c. Propranolol
- d. Lidocaine

118-A patient complaining of changing of his urine color to red/ orange. Which one of the following drugs might cause it?

- a. Augmentin
- b. Dapson
- c. Rifampicin*
- d. Diphenhydramine

119-Misoprostol is classified by FD A as pregnancy category X. that means:

- a. It is absolutely contraindicated during pregnancy*
- b. can be used with caution
- c. Absolutely safe
- d. can be used in the third trimester only

120-Sedating antihistamines effect is related to:

- a. H2 antagonists
- b. H1 antagonists*
- c. A and B
- d. None of the above

121-Intraosseus injection (IO) refers to injection into the:

- a. Bone marrow*
- b. Superficial skin layer
- c. Muscle mass
- d. Pleural fluid

122-The Generic name of Cotrimozazole is:

- a. Bactrim
- b. Ultracef
- c. Comazole
- d. None of the above*

123-There commended dose of paracetamol for a 10 kilogram child is:

- a. 50mg every 6 hours
- b. 150-300mg four times daily
- c. 15mg every 4 - 6hours
- d. 100-150mg every 4-6hours*

124-The PH for acidic preparation is:

- a. 7
- b. More than 7
- b. Less than 7*
- c. 5.5-8.5

125-Eight fluid ounce equal to:

- a. 240ml*(1US fluid ounce = 29.5735)
- b. 120ml
- c. 60ml
- d. none of the above

126-Which of the following drugs should not be mixed with normal saline for intravenous administration

- a. Phenytoin
- b. Cefazolin
- b. Amphotericin B*
- d. Insulin

127-Which of the following drugs should not be mixed with dextrose for intravenous administration

- a. Phenytoin*
- b. Cefazolin
- b. Amphotericin B
- d. Insulin *

128-Quinolones are NOT recommended in children because:

- a. They are not very effective unlike the adults
- b. They have serious Side effect on liver
- c. They may affect their joints*
- d. They make the child hyperactive

129-Which of the following medications cause Rhabdomyolysis is?

- a. Simvastatin*
- b. Sumatriptan
- c. Phenytoin
- d. Fusidic acid injection

130-Pharmacodynamic is defined as:

- a. The study of the biochemical and physiological effects of the drug.
- b. Drug interactions, dosage form, and Side effects of the drugs.
- c. The study of time course of a drug and its metabolite in the body after administration
- d. All of the above.

131-A father seeks your advice for his 2 months old child complaining of runny nose and Temperature of 39°C. What is your advice?

- a. Give him paracetamol and Actifed.
- b. Watch the child for the next 48 hours.
- c. Take the child to the emergency department. *
- d. Start him immediately on Amoxil.

132-Upper respiratory tract infection are treated sometimes with antibiotics. The comments microorganisms for which you suggest an antibiotic are:

- a. Gram negative organisms.
- b. Gram positive organisms.
- c. Mixed micro organisms. *
- d. None of the above.

133-Histoplasmosis is caused by an internal invasion of the body by:

- a. Virus.
- b. Fungal infection*
- c. Helminthes.
- d. Protozoa. **Is a parasitic disease caused by the protozoan Toxoplasma gondii**

134-The drug of choice for head lice is:

- a. Topical steroids.
- b. Pyrethrins. (Pyrethroids) & Malathion*
- c. Antihistamines.
- d. Antibiotics.

135- All of the following consider Calcium channel blockers EXCEPT:

- a. Felodipine.
- b. Diltiazem.
- c. Nifedipine.
- d. Lamivudine* **potent nucleoside analog reverse transcriptase inhibitor (nRTI).For HIV**

136-Norepinephrine stimulates β_1 adrenergic receptors and α adrenergic receptors causing:

- a. increased contractility and heartrate.
- b. Vasoconstriction and increasing systemic blood pressure and coronary blood flow.
- c. A and b*
- d. None of the above.

137-These drugs cannot be dispensed to patients without a prescription EXCEPT:

- a. Bisacodyl.*
- b. Tegretol.
- c. Pravastatin.
- d. Metformin.

138-The well-known complexation reaction between Warfarin and certain drugs containing metal ions such as aluminum hydroxide leads to:

- a. Formation of complexes in the GI tract that is poorly absorbed.
- b. Formation of complexes in the GI tract that is highly absorbed.
- c. Formation of complexes in the GI tract cause irritation.
- d. None of the above.

Not affected by aluminium hydroxide. Magnesium hydroxide & trisilicate increase absorption

139-Which of the following are secreted from the pancreas?

- a. Insulin +FSH.
- b. Insulin +Bileacids.
- c. Insulin +Glucagon. *
- d. Insulin only

140-The Insulin that cannot be given intravenously is:

- a. NPH.*
- b. Regular.
- c. Insulin Lispro.
- d. All of the above.

141-All the secases not cure with thiazides except:

- a. hypernatrimia.*
- b. hyperglycemia.
- c. hypokalaemia.
- d. hyperuricaemia.

142-How much magnesium sulfate is in 10ml of a 200mg/ml magnesium sulfate solution

- a. 1gm
- b. 2gm*
- c. 1000gm
- d. 2000gm

143-How many milliliters of ampicillin do you have to dispense if the patient needs to take 2tsp qid for 7days:

- a. 280mls*
- b. 560mls
- c. 840mls
- d. None of the above

144- A 20-kg child receives erythromycin 30mg /kg q 6h, so he receives.....mg/day

- a. 180
- b. 3600
- c. 2400*
- d. 120

145-three grains equal.....mg

- a. 168
- b. 65
- c. 195*
- d. None of the above

146-Calcitriol is:

- a. 1,25 Dihydroxy cholecalciferol*
- b. 1 Dihydroxy cholecalciferol
- c. 25 Dihydroxy cholecalciferol
- d. None of the above

147- 25 oC=.....oF

- a. 31.66
- b. 77*
- c. 13
- d. 3.8

$$F=(oC*1.8)+32$$

148-D10W means.....

- a. solution containing 10mg dextrose in 100ml water
- b. solution containing 10mg dextrose in 100L water
- c. solution containing 10gm dextrose in 1000ml water
- d. solution containing 10gm dextrose in 100ml water*

149- 0.5 NS solution contains.....milligram sodium chloride in 30ml.

- a. 0.27
- b. 270
- c. 135
- d. 0.135

150- 1milliliter=.....minims

- a. 14.23
- b. 15.23
- c. 16.23*
- d. None of the above



151- 1deciliter=.....liter

- a. 10
- b. 100
- c. 0.1*
- d. 0.01

152-The concentration of one drug is 1:100,if the patient takes 900ml at the morning and 600 ml in the evening, How much drug will the patient take in15 days?

- a. 15mg
- b. 20mg
- c. 30mg
- d. 40mg *

153-How many milliliters of a NS solution can be made from 4500mg of sodium chloride?

- a. 0.5L*
- b. 5L
- c. 50L
- d. None of the above

154-When the patient want to take Viagra, you should ask him:

- a. If he is asthmatic
- b. His age
- c. If he is taking nitroglycerine*
- d. All of the above

155-How many grams of water need edtoprepare150gm of potassium acetate solution with concentration 4% w/w?

- a. 157
- b. 144*
- c. 66
- d. 6

156-What is the concentration of solution,if we dilute100ml 5% of antiseptic solution to 200ml?

- a. 10%
- b. 5%
- c. 2.5%*
- d. 0.5%

157-A patient who weighs 50kg receives 5mg/kg of acyclovir tid, sothepatient receives.....mg/day

- a. 15
- b. 250
- c. 500
- d. 750*

158-The main neurochemical transmitter in the parasympathetic nervous system is

- a. acetylcholine*
- b. epinephrine
- c. ephedrine
- d. amphetamine

159-Which one of the following is considered non selective β blocker

- a. esmolol
- b. betaxolol
- c. propranolol*
- d. metoprolol

160-Which one of the following β blockers used in ophthalmic drops for treatment glaucoma

- a. esmolol
- b. propranolol
- c. timolol*
- d. sotalol

161-Which one of the following β blockers used as antiarrhythmia as it can block K (potassium) channel

- a. esmolol
- b. sotalol*
- c. timolol
- d. metoprolol

162-Which one of is used as β_2 agonist?

- a. Salbutamol *
- b. butoxamine
- c. clonidine
- d. phenylephrine

163-is oproterenol is sympathetic agonist and can be used by:

- a. injection
- b. inhalation
- c. topically
- d. rectally

164-Which one of the following used as vasodilator and can be used in benign prostatic hypertrophy?

- a. tolazocin
- b. prazosin*
- c. phentolamine
- d. phenoxybenzamine

165-The trade name of prazosin is:

- a. Minipress®*
- b. Flagicure®
- c. Canesten®
- d. Zodium®

166-Which one of the following agents can be used in migraine prophylaxis?

- a. ergotamine
- b. ergonovine
- c. methylergonovine
- d. methysergide*

167-Which one of the following agents can be used as power fuloxytotic?

- a. ergotamine*
- b. ergonovine (ergometrine)
- c. methylergonovine
- d. methsyrgide

168-Which one of the following agents used to Decrease postpartum hemorrhage?

- a. ergotamine
- b. ergonovine
- c. methylergonovine (Methylergometrine)*
- d. methsyrgide

169-Which one of the following agents used for migraine treatment (acute attack)?

- a. ergotamine*
- b. ergonovine
- c. methylergonovine
- d. methsyrgide

170-The anticoagulant effect of warfarin Decreased by these of:

- a. Rifampicin
- b. Aspirin
- c. Vitamin K*
- d. verapamil

171-Which one of the following parasympathomimetic can be used as antiemetic?

- a. carbachol
- b. pilocarpine
- c. bethanecol
- d. metoclopramide*

172-Which one of the following can be used intreatment the toxicity of organophosphorus compound?

- a. ecothiopate
- b. pyridostigmine
- c. parathione
- d. parlidoxime*

173-candidiasis is considered

- a. viralinfection
- b. bacterialinfection
- c. fungalinfection*
- d. protozoainfection

174-histoplasmosis is considered

- a. viralinfection
- b. bacterial infection
- c. fungal infection*
- d. protozoa infection

175-amebiasis or amoebic dysentery can be treated by

- a. metronidazole*
- b. flubendazole
- c. alpendazole
- d. nifurtimox

176-Which one of the following parasymphathetic agent can be used as ophthalmic drops to treat glaucoma?

- a. pilocarpine*
- b. bethanecol
- c. metoclopramide
- d. edrophonium

177- All of the following are atropine Side effects except:

- a. constipation
- b. diarrhea*
- c. drymouth
- d. rapidpulse

178-Which one of the following is considered sedating antihistaminic?

- a. loratidine
- b. astemizole
- c. diphenhydramine*
- d. acrivastine

179-Claritin® is a trade name of a drug and the Generic name is?

- a. loratidine*
- b. astemizole
- c. diphenhydramine
- d. acrivastine

180-fexofenadine is antiallergic and can be sold under the name :

- a. Trictine®
- b. Telfast®*
- c. Allergex®
- d. Pirafen®

181-The H1 blocker is used as:

- a. antiallergic*
- b. ulcer
- c. antidiarrhea
- d. antiamobeic

182-The H2 blocker is used as:

- a. antiallergic
- b. Ulcer*
- c. antidiarrhea
- d. antiamobeic

183-Which one of the following used for treatment peptic and duodenal ulcer?

- a. astemizole
- b. cimetidine*
- c. ticlopidine
- d. acrivastine

184-The trade of ranitidine is KSA is:

- a. Histac®
- b. Zantac®*
- c. Tagamet®
- d. Anafranil®

185-The trade name of cimitidine is:

- a. Histac®
- b. Zantac®
- c. Tagamet®*
- d. Anafranil®

186-Benzodiazepine is preferred as sedative hypnotic than barbiturate:

- a. as benzodiazepineis more safer
- b. as benzodiazepineis less liable to produce tolerance
- c. barbiturate is more liable to produce tolerance
- d. All of the above*

187-Which one of the following is considered ultra short acting barbiturate?

- a. thiopental*
- b. secobarbital
- c. amobarbital
- d. phenobarbital

188-Which one of the following is considered short acting barbiturate?

- a. thiopental
- b. secobarbital*
- c. amobarbital
- d. phenobarbital

189-Which one of the following is considered intermediate acting barbiturate?

- a. thiopental
- b. secobarbital
- c. amobarbital*
- d. phenobarbital

190-Which one of the following is considered long acting barbiturate?

- a. thiopental
- b. secobarbital
- c. amobarbital
- d. phenobarbital*

191-Leprosy can be treated by

- a. isoniazide plus streptomycin
- b. dapsone and clofazimine*
- c. rifampin and streptomycin
- d. rifampin and ethambutol

192-Xanax® is sedative drugs and the Generic name is:

- a. diazepam
- b. chlorazepam
- c. alprazolam*
- d. clorazepate

193-Which one of the following can be for treatment benzodiazepine over dose:

- a. disulfiram
- b. buspirone
- c. flumazenil*
- d. parlidoxime

194-Which one of the following can be used for treatment alcohol addiction?

- a. disulfiram*
- b. buspirone
- c. flumazenil
- d. parlidoxime

195-Which one of the following can be as anxiolytic drug?

- a. disulfiram
- b. buspirone*
- c. flumazenil
- d. parlidoxime

196-Which one of the following can be used in morphine over dose?

- a. methadone
- b. naloxone*
- c. pethidine
- d. alfentanil

197-Death can occur in morphine toxicity due to:

- a. Hypotension
- b. Respiratory failure*
- c. Cardiac arrest
- d. Meningitis

198- All of the following can be used as antitussive except:

- a. codeine
- b. dextromethorphan
- c. metronidazole*
- d. phloco diene

199-Cholera can be treated by

- a. doxacycline*
- b. penicillin
- c. cephalosporin
- d. none of the above

200-The death occur in cholera due to

- a. respiratory failure
- b. Cardiac arrest
- c. Dehydration*
- d. none of the above

201-T.B is abbreviation for tuberculosis which can be treated by

- a. isoniazide,rifampin,ethambutol & pyrazinamide*
- b. isoniazide,rifampin,ethambutol & chloramephenicol
- c. isoniazide,rifampin,ethambutol & dapsone
- d. isoniazide,rifampin,ethambutol & clofazimine

202-Flagyl® is at trade name of:

- a. metronidazole*
- b. flubendazole
- c. alpendazole
- d. nifurtimox

203-The active ingredient in Leponex® is:

- a. Clozapine*
- b. Respiredone
- c. Lithium
- d. Haldol

204- All of the following are considered selective serotonin reuptake inhibitor (SSRIs) except:

- a. Fluoxetine
- b. Citalopram
- c. Sertraline
- d. Imipramine*(TCA)

205-Parkinsonism can be treated by All of the following except:

- a. L-dopa
- b. Dopamine*
- c. amantadine
- d. bromocriptine

206-Which one of the following is used for treatment Parkinsonism?

- a. Sinemet®*
- b. Stellazil®
- c. Stellazine® **Trifluoperazine**
- d. Stroka®

207-carbamazepine is antiepilepsydrugs oldunderthetrade name :

- a. Bional®
- b. Taural®
- c. Epiran®
- d. Tegretol®*

208-Status epilepticus best treated by:

- a. diazepam*
- b. valproicacid
- c. phenytoin
- d. Primidone

209-ethusoxamide can be used for treatment of :

- a. petitmalepilepsy*
- b. grandmalepilepsy
- c. statusepilepticus
- d. partial status

210-phenytoinisGeneric name of:

- a. Epanutin®*
- b. Cidoquin®
- c. Dusputaline®
- d. Idiavix®

211-Pseudomembranous colitis can be treated by

- a. vancomycin plus metronidazole*
- b. vancomycin plus tetracycline
- c. ampicillin plus metronidazole
- e. amoxacyillin plus metronidazole

212-Gonorrhoea is infectious disease and can be transmitted

- a. sexually*
- b. topically
- c. Inhalation
- d. All of the above

213-Avianflu(Birdflu) can be treated by:

- a. zidovidine
- b. amantadine
- c. ganciclovir
- d. oseltamivir*

214-all these drugs are antibacterial except:

- a. zalcitabine*
- b. lomefloxacin
- c. cefopyrazone
- d. Nalidixic acid

215-Which of the following antibiotics can be used topically?

- Acyclovir.
- Cefaclor.
- Cephalexin.
- Fusidicacid.*

216-all these are natural Estrogens except:

- a. Mestronol*
- b. Oestrone
- c. Oestriol
- d. Oestradiol

217-Which of the following has vasodilator effect?

- a. nicotine*
- b. cholestyramine
- c. urokinase
- d. vit.A

218-One of the following is not from Quinolone group:

- a. naldixic acid
- b. quinine*
- c. eoxacin
- d. norfloxacin

219-One of the following is not for epilepsy:

- a. clorizepam
- b. phenytoin
- c. primidone
- d. imipramine*

220- All of these can be used for leprosy except:

- a. dapson
- b. rifampicin
- c. clofazimine
- d. INH*

221-Sumatriptan is used for:

- a. Hypertension
- b. treatment of migraine*
- c. angina pectoris
- d. control of epilepsy

222-Which is enzyme inducer?

- a. rifampicin*
- b. cimetidin
- c. vit.C
- d. chloramephenicol

223- All of the following for prophylaxis of asthma except:

- a. Ketotifen
- b. nedocromil sod.
- c. Sodium cromoglycate
- d. Salbutamol *

224- All of the following is Non Sedating Anti-Histamine, except:

- a. Loratadine
- b. Cetrizine
- c. Astimazole
- d. Azotodin*

225-Herpes simplex can be treated by

- a. ganciclovir
- b. acyclovir*
- c. zalcitabine
- d. oseltamivir

226- All of the following are controlled drugs except:

- a. rivotril
- b. epanutin*
- c. sresolid
- d. diazepam

226-all these drugs are anti-viral except:

- a. amantadine
- b. zidovudine
- c. acyclovir
- d. aluerin*(antispasmodic)

227-Which one of these symptoms not occur with morphine?

- a. diarrhea*
- b. respiratory depression
- c. constipation
- d. vomiting

228-Which one of these drugs not used for acute pain?

- a. naproxen
- b. colchicine
- c. codeine
- d. prednisolone*

229-The Recommended daily dose of Vitamin A for correction of deficiency is:

- a. 5,000IU
- b. 10,000IU
- c. 20,000IU
- d. 30,000IU*

230-The daily dose of Vitamin A for therapy is:

- a. 10,000IU
- b. 20,000IU
- c. 50,000IU
- d. 100,000IU*

231-Hepatitis vaccine dose is:

- a. once/year
- b. twice/year
- c. three times /year*
- d. four times /year

232-Which one is OTC drug?

- a. Simple analgesic*
- b. Warfarin
- c. Ciprofloxacin
- d. Amoxicillin

233-One of the following is given once daily:

- a. Amorphous Insulin
- b. Regular Insulin
- c. Neutral Insulin
- d. Protamine zinc Insulin *

234-all these are anticoagulants except:

- a. warfarin
- b. ancord
- c. heparin
- d. dipyridamol*

235-all these are anti-platelets except:

- a. dipyridamol
- b. aspirin
- c. streptokinase*
- d. ticlopidine

236-Acyclovirisa Generic name soldin KSA under the name

- a. Trosyd®
- b. Mycospor®
- c. Zovirax®*
- d. Amikacin®

237-Ferrograde is:

- a. ferrous sulphate*
- b. ferrous citrate
- c. ferrous gluconate
- d. ferrous fumarate

238-Procaïnamide is:

- a. Nachannel blocker *
- b. nifedipine
- c. lisinopril
- d. quinidine

239-Thiazide diuretics cause all the following except:

- a. hypokalaemia
- b. hypercalcemia
- c. hyperuricaemia
- d. hypernatremia*

240-Elimination of digoxin is mainly by:

- a. kidney*
- b. liver
- c. both
- d. none of them

241-What is the most drug that can cause damage if it is given OTC?

- a. captopril
- b. warfarin*
- c. chlorothiazide
- d. rifampicin

242-All are antimetabolites except:

- a. methotrexate
- b. flurouracil
- c. cytarabine
- d. cyclophosphamide*
- e. azathioprine(Imuran)

243-Cefaclor is:

- a. 1st generation cephalosporin
- b. 2nd generation cephalosporin*
- c. 3rd generation cephalosporin
- d. 4th generation cephalosporin

244-Cefixime is:

- a. 1st generation cephalosporin
- b. 2nd generation cephalosporin
- c. 3rd generation cephalosporin*
- d. 4th generation cephalosporin

245-Zinnat is:

- a. 1st generation cephalosporin
- b. 2nd generation cephalosporin*
- c. 3rd generation cephalosporin
- d. 4th generation cephalosporin

246-Which aminoglycoside antibiotic can be taken orally?

- a. gentamycin
- b. neomycin*
- c. amikacin
- d. tobramycin

247-all these medication sare used intreatment of TB except:

- a. cyclosporin*
- b. INH
- c. cycloserin
- d. rifampicin

248-One of these is not like others:

- a. timolol
- b. acebutalol
- c. labetolol*
- d. propranolol

249-Betahistine HCl (betaserc) used for:

- a. formieniere`s disease*
- b. antihistamine
- c. hypersensitivity reactions.
- d. All of the above.

250-Which one is not 5HT3 antagonist?

- a. tropisetron
- b. ondansetron
- c. domperidone*
- d. gransetron

251-Which is not oral anti-coagulant?

- a. warfarin
- b. enoxaprine*
- c. phenidione
- d. nicoumolone

252- All of these are the components of calamine lotion B.P except:

- a. sod.Sulphate*
- b. calamine
- c. sod.citrate
- d. bentonite
- e. Zinc Oxide

253-Terbinafene is used topically for:

- a. eczema
- b. candida
- c. fungal infections*
- d. antibiotic

254-Which one is used for benign prostatic hyperplasia?

- a. flumazenil
- b. cefprozil
- c. mivacurim
- d. finasteride*

255- All of these are 3rd generation cephalosporin except:

- a. cefixime
- b. cefopodoxime
- c. cefotriaxone
- d. cefotaxime

256-Which of the following is not mucolytic?

- a. bromohexin
- b. dextromethorphan*
- c. ambroxol
- d. guaphenesin

257-Which one is anti-platelet?

- a. ticlopidine*
- b. desfluron
- c. aldesleukin
- d. atovouon

258- All of these are anti-neoplastic except?

- a. aldesleukin
- b. taniposide
- c. pentostatin
- d. amlodipine*

259-Which is not benzodiazepne?

- a. temazepam
- b. nitrozepam
- c. loprazepam
- d. clozapine*

260-Oralre hydrate solution(ORS) contains:

- a. sod.citrate, pot.chloride & glucose
- b. sod.citrate, pot.chloride & sod.chloride
- c. sod.chloride, pot.citrate, sod.citrate&glucose*
- d. sod.chloride, pot.chloride, sod.bicarbonate & citricacid.

261-Which of the following is selective α_1 blocker ?

- a. labitolol
- b. terazosine*
- c. clonidine
- d. captopril
- e. verapamil

262-Allopurinol is used as:

- a. analgesic agent
- b. uricosuric agent*
- c. antiinflammatory agent
- d. antipyretic agent

263-Which of the following is released by bacterial infection?

- a. endotoxin*
- b. exotoxin
- c. antibiotics
- d. cytotoxin

264-The amount of water in adult male is about:

- a. 25%
- b. 60%*
- c. 80%
- d. 10%

265-Which of the following is the first choice in acute gout?

- a. allopurinol
- b. indomethacin*
- c. colchicine

266-Which of the following is NOT a risk factor for osteoporosis?

- a. Minimal exercise
- b. Low calcium intake
- c. Male gender*
- d. Family history

267-The latin abbreviation for “ After Meals” is:

- a. a.c.
- b. a.a.
- c. p.c.*
- d. i.c.
- e. c.c

268-The latin abbreviation for “Every Night” is:

- a. a.c.
- b. o.n.*
- c. p.c.
- d. i.c.
- e. c.c

269-Which of the following consider Parenteral administration ?

- a. Subcutaneous*
- b. oral
- c. aa
- d. all

270-INH can be used for treatment of TB.

- a. true*
- b. False

271-Fentanyl is considered controlled medication:

- a. true*
- b. False

272-Rosiglitazon is useful in treatment of :

- a. DM*
- b. HF
- c. CKD
- d. HTN

273- 8gm of water are needed to make 160gm of 5%w/w solution of K acetate.

- a. true
- b. False*

274-Which of the following are given IV rout?

- a. Atrovastatin
- b. Gentamycin
- c. Nitroglycerin
- d. b,c*

275-The main neurochemical transmitter in the parasympathetic nervous system is

- a. acetylcholine*
- b. Epinephrine
- c. Ephedrine
- d. Amphetamine

276-Which one of the following is considered non selective β blocker

- a. esmolol
- b. betaxolol
- c. propranolol*
- d. metoprolol

277--Which one of the following β blockers used in ophthalmic drops for treatment glaucoma

- a. esmolol
- b. propranolol
- c. timolol*
- d. sotalol

278-Which one of the following β blockers used as anti arrythmia as it can block K (potassium) channel

- a. esmolol
- b. sotalol*
- c. Timolol
- d. Metoprolol

279-Which one of the following is used as β_2 agonist?

- a. Salbutamol *
- b. Butoxamine
- c. clonidine
- d. Phenylephrine

280-Which one of the following used as vasodilator and can be used in benign prostatic hypertrophy

- a. tolazocin
- b. prazosin*
- c. Phentolamine
- d. Phenoxybenzamine

281-Which one of the following agents can be used as powerful oxytocic?

- a. ergotamine*
- b. ergonovine (ergometrine)
- c. methylergonovine
- d. Methsyrgide

282-Which one of the following drugs doesn't make the feces black

- a. ferric gluconate
- b. ferroic
- c. rifampicin
- d. warfarine
- e. phenytoin*

283-Which one of the following leads to constipation

- a. aspirin
- b. ibuprofen
- c. ca carbonate
- d. aluminium hydroxide*
- e. morphine*

284- 10 grains equal how much grams

- a. 0.0065
- b. .065
- c. .65*(1g=15gr)
- d. 6.5
- e. 65

285-The mixture of liquid paraffine and water will be

- a. solution
- b. lotion
- c. emulsion*
- d. elixir
- e. drops

286-Which one of the following not be fibrinolytic drug

- a. streptokinase
- b. urokinase
- c. alteplase
- d. heparine*

287-Types on nerves in the nervous system

- a. motor
- b. Sensory
- c. Mixed
- d. a and b
- e. All of the above*

288-Which one of the following shows anti pyretic activity?

- a. amiodorone
- b. metrinidazol
- c. mebendazol
- d. Nitroglycrine
- e. diclofenacNa*

289-Roacutene (isotretinoin) can be used for

- a. acnevulgaris*
- b. TB
- c. Amoeba
- d. Hypertension
- e. DM

290-Roacutene is

- a. Catecholamine
- b. Vitamin A derivative*
- c. Cardiacglycoside
- d. Alkaloids
- e. Vitamin D drevatives

291-which of the following sentences is true

- a. aprazolam should not be used for longterm*
- b. Aspirin is safe in peptic ulcer patient
- c. Ciprofloxacinis safe during pregnancy
- d. Morphineca use diarrhea
- e. Atropine indicated in Glaucoma

292-Do butamine mustn't be taken

- a. IV
- b. Orally*
- c. IM
- d. SC
- e. All above

293-Prednisolone used for treatment

- a. Addison's disease
- b. Hyponatremia
- c. Severe bronchial asthma
- d. All of the above*
- e. None of the above

294-Which one of the following is not pure β - blocker

- a. betaxolol
- b. sotalol
- c. propranolol
- d. Labetalol*
- e. Timolol

295-Tirofiban is

- a. prevent blood clotting*(antiplatelet)
- b. Prevent fungal growth
- c. Antibiotic
- d. NSAID
- e. ACEI

296-Which one of the following is not considered endocrine gland

- a. Pituitarygland
- b. Adrenalgland
- c. Thyroidgland
- d. Sweatglands*
- e. Ovary

297-Which one of the following is α a gonist?

- a. Phentolamine
- b. Dobutamin
- c. Phenylephrin*
- d. Pilocarbine
- e. Serotonin

298-Which of the following sentences is true about carvedilol

- a. It's Parasympathomimetic
- b. α and β blocker *
- c. β 1
- d. β 2
- e. All above

299-The function of the thyroid gland is/are

- a. growth
- b. memory
- c. body temperature
- d. energy production
- e. All above *

300-Which one of the following has vasodilator effect

- a. Verapamil
- b. Digoxin
- c. Cimetidine
- d. Hydralazine*
- e. Fluoxetine

301-Symmetrel® is a trade name used for treatment

- a. Diabetes
- b. Parkinsonism*(amantadine)
- c. Peptic Ulcer
- d. Fever
- e. Cancer

302-Which one of the following drugs can be used for treatment Parkinson's disease?

- a. Rantidine
- b. Amlodipine
- c. Prazosine
- d. Bromocriptine*
- e. Phenobarbital

303-oxytocin is category X so it

- a. shouldn't be given during pregnancy*
- b. Safe during pregnancy
- c. Given with caution
- d. Risk is unknown
- e. non above

304-Which one of the following drugs shouldn't be taken during pregnancy

- a. enalapril*
- b. sucralfat
- c. ibuprofen
- d. paracetamol
- e. aldomet

305-the therapeutic index is refer to relation which is

- a. (-)
- b. (+)
- c. (X)
- d. (/)*

306- Insulin shouldn't be used with

- a. dextrose*
- b. Saline
- c. Mannitol
- d. Aspirin
- e. Paracetamol

307-Which one of the following is not Insulin Side effect ?

- a. atrophy at site of injection
- b. hypoglycemia
- c. hyperglycemia*
- d. all are Side effect
- e. all are not Side effect

308-Which one of the following is considered Glimipride Side effect

- a. hypoglycemia*
- b. hyperglycemia
- c. all are Side effect
- d. all are not Side effect

309-Which one of the following drugs can't be taken orally?

- a. Enoxparine*
- b. Ampicillin
- c. Aspirin
- d. neomycin
- e. Atropin

310-Which one of the following is not cephalosporin?

- a. clarithromycin*
- b. Cephalexin
- c. Cefoxitin
- d. Cephmandole
- e. Cefepime

311-The type II Diabetes Mellitus can be treated with:

- a. sulphonylurea
- b. exercise
- c. Insulin
- d. Metronidazol
- e. a and b*

312-Which one of the following is not OTC drugs

- a. Bisacodyl
- b. Aspirin
- c. Duphlac
- d. Brufen
- e. warfarine*

313-Which of the following symptoms is due to increase epinephrine in the sympathetic nervous system:

- a. Increase Heartrate
- b. Bronchodilator
- c. Mydriasis
- d. Vasoconstriction
- e. All of the above*

314-The mixture of acacia, water, and oil will be

- a. lotion
- b. Suspension
- c. Solution
- d. Emulsion*

315-ondansetron HCl is used for

- a. Hypertension
- b. angina pectoris
- c. prophylaxis of migraine
- d. anti-emetic*

316-all these are Side effects of chloramphenicol except

- a. irreversible aplastic anemia
- b. opticneuritis
- c. peripheral neuritis
- d. iron deficiency anemia*

317-ticlopidine is used for

- a. anticoagulant
- b. antibiotic
- c. anticholinergic
- d. antiplatelets*

318-all these drugs are fibrinolytic except

- a. Dipyridamole*
- b. streptokinase
- c. urikinase

319-all these drugs can be used for parkinsonism except

- a. selgiline HCl
- b. carbidopa
- c. pergolide
- d. nedocromil sodium*

320-cardiac arrest could be managed by the following except

- a. ephedrine
- b. lidocaine
- c propranolol*
- d. normal saline

321- Insulin which can be given I.V. is

- a. regular type*
- b. 30/70 type
- c. NPH type
- d. None of the above

322-Drug used for leishmania is

- a. sodium stiboglyconate*
- b. sodium cromoglyconate
- c. mebendazole

323-all these drugs are antibacterial except

- a. Zalcitabine*
- b. lomefloxacin
- c. cefoprazone

324-all these are natural estrogens except

- a. Mestronol*
- b. oestrone
- c. oestriol
- d. oestradiol

325-co-trimethoxazole contains

- a. trimethoprim + sulphamethoxazole*
- b. trimethoprim + sulphisoxazole
- c. trimethoprim + sulphadiazine
- d. trimethoprim + sulphacetamide

326-Which one of these drugs is the drug of choice for myxedema

[hypothyroidism]

- a. Carbimazole*
- b. iodine
- c. thyroxinesodium
- d. propylthiouracil

327-benzyl peroxide is

- a. local treatment of acnevulgaris*
- b. local treatment of scabies
- c. local treatment of inflammation
- d. local treatment of pain

328-thiazide diuretics cause all the following except

- a. hypokalemia
- b. hypercalcemia
- c. hyperuricemia
- d. hyponatremia
- e. hyperglycemia
- f. hypernatremia*

329-treatment with anti-arrythmic drug depends on except

- a. type of medication
- b. period of medication
- c. patient sex*

330-amilorid as trimetren causes

- a. CNS disturbances
- b. Hyperkalemia*
- c. Hypokalemia
- d. Not affect potassium

331-Which one is not tetracycline

- a. minocycline
- b. demeclocycline
- c. doxycycline
- d. clindamycin*

332-Which one is not Calcium channel blocker

- a. verapamil
- b. diltiazem
- c. amlodipine
- d. cinnarzine*

333-all these drugs are used intreating T.B. except

- a. Cyclosporine*
- b. INH
- c. Cycloserin
- d. Rifampin

334-all these drugs are anti-cholinergic except

- a. iprotropium
- b. atropine sulphate
- c. benztropine methylate
- d. dobutamin*

335-which is correct about diazepam

- a. can cause muscle spasm
- b. is taken I.M. only
- c. is derivative from morphine
- d. is not used forlongterm*

336-maximum dose of paracetamol is

- a. 4gm/day*
- b. 5gm/day
- c. 7gm/day
- d. 10gm/day

337-Metoclopramide is

- a. dopamine antagonist*
- b. acts centrally by blocking vomiting center
- c. acts peripherally on the chemoreceptor trigger zone (CTZ)
- d. used as anti emetic*
- e. All of the above

338-Which one is contraindicated in pregnancy

- a. sucralfate
- b. ethambutol
- c. chlorpheniramine
- d. ciprofloxacin*

339-the antidote for heparin is

- a. protamine sulphate*
- b. acetyl cysteine
- c. Vitamin K
- d. naloxone

340-the antidote for paracetamol

- a. Protamine sulphate
- b. acetyl cysteine*
- c. Vitamin K
- d. naloxone

341-what is the Generic name of vitamin B2

- a. cyanocobalamin
- b. niacin
- c. riboflavin*
- d. pantothenic acid
- e. pyridoxine
- f. folic acid

342-what is the Generic name of vitamin B3

- a. cyanocobalamin
- b. niacin*
- c. riboflavin
- d. pantothenic acid
- e. pyridoxine
- f. folic acid

343-what is the Generic name of vitamin B5

- a. cyanocobalamin
- b. niacin
- c. riboflavin
- d. pantothenic acid*
- e. pyridoxine folic acid

344-what is the Generic name of vitamin B6

- a. cyanocobalamin
- b. niacin
- c. riboflavin
- d. pantothenic acid
- e. pyridoxine*
- f. folicacid

345-what is the Generic name of vitamin B9

- a. cyanocobalamin
- b. niacin
- c. riboflavin
- d. pantothenic acid
- e. pyridoxine
- f. folicacid*

346-what is the Generic name of vitamin B12

- a. cyanocobalamin*
- b. niacin
- c. riboflavin
- d. pantothenic acid
- e. pyridoxine
- f. folicacid

347-what is the Generic name of vitamin B1

- a. cyanocobalamin
- b. niacin
- c. riboflavin
- d. Thiamin*
- e. pyridoxine
- f. folicacid

348-fluxotine is

- a. prozac
- b. selective serotonin reuptake inhibitor
- c. anti inflammatory drug
- d. a & b*

349-omeprazole is

- a. losec
- b. proton pump inhibitor
- c. analgesic
- d. a & b*

350-which of the following diseases that cause edema

- a. CHF
- b. Renal failure
- c. Arrhythmia
- d. a & b*

351- All of these are the components of calamine lotion B.P. except

- a. ZnO
- b. Calamine
- c. Sodiumcitrate
- d. Bentonite
- e. Sodium sulphate*

352-MMR vaccine is

- a. measles vaccine
- b. mumps vaccine
- c. rubella vaccine
- d. measles-mumps-rubella vaccine*

353-sodium nitroprusside

- a. potent vasodilator*
- b. potent diuretic
- c. potent vasoconstrictor
- d. anti-arrhythmicagent

354-a drug needs monitor for its serum level

- a. amikacin
- b. warfarin
- c. theophyllin
- d. digoxin
- e. All of the above*

355-for fluxacillin, all these statements are wrong except

- a. it is resistant to penicillinase enzyme
- b. it is taken by oral route only*
- c. it is base stable it is better absorbed than cloxacillin

356-thiabendazole is one of thiazide diuretics

- a. true
- b. false*

(it is an anthelmintic)

357-praziquantil is

- a. causes loss of intracellular calcium of worms
- b. used for malaria
- c. biltracidis the trade name *
- d. a & c

(↑Caentry)

358-nitrofurantoin is indicated for

- a. viral infection
- b. GIT infection
- c. Urinarytract infection*
- d. URI

359-Vitamin K

- a. increase formation of clotting factor*
- b. increase conversion of vit.K to the active form
- c. prevent conversion of Vitamin K back to the active form
- d. none of the above

360-quinidine sulphate is

- a. anti-arrhythmic agent
- b. antimalarial agent
- c. antihypertensive agent
- d. a & c*

361-ACE stands for

- a. converting Ag II to vasoconstrictor Ag I
- b. converting Ag I to vasoconstrictor Ag II*
- c. converting AgI to vasodilator Ag I
- d. converting Ag II to vasodilator Ag I

362-treatment of anaphylactic reaction is initiated by immediate

- a. Hydrocortisone
- b. cimetidine
- c. human albumin
- d. epinephrine*

363-introduction of emesis is not indicated in the following except

- a. caustic substance ingestion
- b. CNS
- c. Seizures
- d. Paracetamol poisoning*

364-the following are non sedating anti-histamines except

- a. loratadine
- b. cetirizine
- c. terfenadine
- d. famotidine*

365-type II diabetics is characterized by

- a. lack of response to Insulin
- b. lack of response to sulphonylurea
- c. respond to dietary therapy alone
- d. slowly increasing sulphonylurea requirement*

366-the following are Side effects of aminoglycosides except

- a. nephrotoxicity
- b. ototoxicity
- c. eyetoxicity*
- d. abdominal pain (N&V)

367-the following are potassium sparing diuretics except

- a. trimetren
- b. spironolactone
- c. amiloride
- d. furosemide*

368-Which one reduces the metabolism of other drugs in the liver

- a. rifampin
- b. vit.C
- c. cimetidine
- d. metronidazole

369-Specific serotonin reuptake inhibitors are used as

- a. antihistaminic
- b. antidepressant*
- c. antihypertensive
- d. antiemetic

370-which of these has no vasodilator effect

- a. Methyldopa*
- b. hydralazine
- c. enalapril
- d. prazosin

371-Calcipotriol is indicated for

- a. psoriasis*
- b. Hypertension
- c. calcium supplement
- d. acne

(vitaminD₃ derivative, induce differentiation and to suppress proliferation of keratinocytes)

372-amantadine is used for All of the following except

- a. antiviral
- b. for influenza A
- c. anti parkinsonism
- d. for influenza B*

373-all are viral infections except

- a. hepatitis B
- b. influenza
- c. herpes zoster
- d. rickettsia*

374-at the cardiac arrest, the drug of choice is

- a. Epinephrine*
- b. propranolol
- c. hydrocortisone
- d. saline solution

375-drugs to pass B.B.B.should be

- a. unionized hydrophilic
- b. ionized hydrophilic
- c. unionized lipophilic*
- d. ionized lipophobic

376- Side effects of Insulin

- a. local reaction
 - b. hypoglycemia
 - c. fat dystrophy at site of injection
 - d. All of the above*
- none of the above

377-one of the following is the main Side effect of Insulin

- a. Hypoglycemia*
- b. fluid retention
- c hyperglycemia
- d. All of the above

378-chloramphenicol affects liver inhibitory enzyme.So

- a. Decrease metabolism & increase response*
- b. Decrease metabolism & Decrease response
- c. increase metabolism & increase response
- d. Decrease metabolism & Decrease response

379-Enteric coated tablets

- a. pass the stomach and release in the intestine*
- b. pass the stomach and intestine
- c. release in the stomach

380- All of these are true for levodopa except

- a. precursor of dopa
- b. does not pass B.B.B*
- c. effective in parkinsonism
- d. stimulate dopamine

381-The conc. Of the drug on expiration date is

- a. not less than 100%
- b. not less than 90%
- c. from 90-110*%
- d. none of the above

better answer:(from 95-105%)

382-New and best therapy for peptic ulcer is

- a. antacid + H2 antagonist*
- b. use more than one H2 antagonist
- c. four medicines including antibiotic

383-all the following change the color of urine except

- a. nitrofurantoin
- b. phenazopyridine
- c. rifampicin
- d. none of the above*

384-which of the following is not cardio selective blocker

- a.atenolol
- b.acebutolol
- c.sotalol
- d.Labetalol*

385-a patient has a prescription of erythromycin 500mg tablets every 6hours and you do not have this antibiotic.You can phone the doctor to change it with

- a. Clarithromycin*
- b. amoxicillin
- c. clindamycin
- d. ampicillin

386-ACE inhibitors are used for the treatment of

- a. asthma
- b. migraine
- c. Hypertension *
- d. All of the above

387-Atropine is used as pre anesthetic to

- a. give skeletal muscle relaxation
- b. Decrease saliva*
- c. Treat the Side effects of anesthetics
- d. Help the sleeping effects of anesthetics

388-disulfiram is used for treatment of

- a. chronic alcohol dependence*
- b. protection of liver cells from alcoholism
- c. prevention of conversion of acetaldehyde to acetic acid
- d. All of the above

389-The following are the indications of diazepam except

- a. antipsychotic
- b. epilepsy
- c. analgesic*
- d. hypnotic

390-Regular use of sublingual doses of Nitroglycerin is likely to result in:

- a. Hepatotoxicity
- b. Nephrotoxicity
- c. Pepticulcer
- d. Tolerance*

391-The tricyclic antidepressant Imipramine (Tofranil) has been approved by FDA for use in:

- a. Parkinson's disease
- b. Hypertension
- c. Enuresis*
- d. Peptic ulcer
- e. None of the above

392- All of the following are beta blockers except:

- a. Metoprolol(Lopressor)
- b. Nadolol(Corgard)i
- c. Enalapril(Renitec)i*
- d. Propranolol(Inderal)i

393-Buspirone (Buspar) is classified pharmacologically as:

- a. Anti depressant
- b. Anxiolytic agent*
- c. Narcotic analgesic
- d. Hypnotic

394-Carbocysteine is an agent used clinically as:

- a. Laxative
- b. Cough suppressant
- c. Diagnostic agent
- d. Mucolytic*

395-treatment of Hypothyroidism:

- a. Cimetidin
- b. Thiamin
- c. Carbimazol
- d. Thyroxine Sodium*

396-treatment of Hyperthyroidism:

- a. Carbimazole
- b. Propylthiouracil
- c. Thyroxine
- d. A and b*

397-Trade name of Nifedipine:

- a. Voltaren
- b. Diamicron
- c. Zantac
- d. Adalat*

398-NSAIDs means:

- a. NO Side effect drugs
- b. Steroid anti inflammatory
- c. Nonsteroid anti inflammatory drugs *
- d. Non symptoms diseases

399-Which of the following not given IV:

- a. Suspension
- b. Emulsion*
- c. Solution
- d. All of the above

400-Short acting barbiturate used as:

- a. Anti depressant
- b. Anesthetic*
- c. Antiepileptic
- d. Diuretics

401-BACITRACIN used :

- a. Only parenteral
- b. Only to pically*
- c. Only orally
- d. All of the above

402-Which of the following are contraindicated with Digoxine:

- a. Aspirin
- b. Spironolactone*
- c. Zocor
- d. Rantidine

403-MEFANAMICACID is:

- a. Anticoagulant
- b. Anti-inflammatory*
- c. Antibiotic
- d. Broncho dilator

404-Which of the following drugs are taken before meal?

- a. VOLTAREN
- b. AUGMENTIN*
- c. GRISOFULVIN (with or after meals)
- d. BRUFEN

405-Antigout:

- a. COLCHICINE
- b. ASPIRIN
- c. ALLOPURINOL*
- d. A and b

406-Treatment of Herps:

- a. NIFIDEPIN
- b. DIGOXIN
- c. ACYCLOVIR*
- d. HYDRALAZIN

407-Maximum dose of Brufen:

- a. 1200mg/day
- b. 3200mg/day*
- c. 4000mg/day
- d. 200mg/day

408-Antiepileptic:

- a. carbamazepin
- b. phyntoin
- c. phenobarbital
- d. All of the above*

409-Scientific name of Vitamin A:

- a. Retinol*
- b. Thiamine
- c. Cyanocopalamin
- d. Calciferol

410-H2 receptor antagonist

- a. Cimetidine*
- b. loratidine
- c. celexocib
- d. Promthazine

411-Which of the following are corticosteroids?

- a. amantadin
- b. hydrocortiesione
- c. prednisoline
- d. b and c*

412-Scientific name of Vitamin K:

- a. Retinol
- b. Phytomenadione*
- c. Niacin
- d. Riboflavin

413-Trade name of Gliclazide:

- a. amoxil
- b. minidiab(glipizide)
- c. diamicon*
- d. Ventolin

414-Generic name of Claritin:

- a. histadin
- b. adrenalin
- c. loratidin*
- d. chloropheniramin

415-Generic name of Buscopan:

- a. atenolol
- b. hyoscine(scopolamine)*
- c. pilocarpine
- d. atropine

416-Trade name of Allopurinol:

- a. profenide
- b. diurecif
- c. voltaren
- d. zyloric*

417-treatment of grand mal epilepsy:

- a. atenolol
- b. phyntoin*
- c. heparin
- d. diclofenac

418-treatment of (petit mal epilepsy):

- a. fluxitin
- b. ethosuximide*
- c. atenolol
- d. warfarin

419-Not used in Dysenteric Amoeba treatment:

- a. Gentamycine*
- b. iodoquinol
- c. metronidazol
- d. diloxanide furoate

420-Dose of Cephalexin:

- a. 1 gm/day
- b. 250mg-500mg/day
- c. 250mg-1000mg/6hr
- d. 500mg/8hr*

421- All of the following are Antiarrhythmic drugs except:

- a. Quinidin
- b. Procainamide
- c. Halothan*
- d. Phyntoine

422-Amino glycosides Mechanism of action:

- a. Inhibit cell wall synthesis
- b. Bind to 30 Sribosome*
- c. Interfere with osmosis characters
- d. Inhibit DNA transfer

423-Drug administered.....are exposed to first-pass effect

- a. orally*
- b. I.M
- c. I.V
- d. S.C

424-Which one these is pro-drug?

- a. ampicillin
- b. erythromycin
- c. levodopa*
- d. All above

425-Bioavailability is defined as:

- a. the action of the body on the drug
- b. rate and extent to which the drug enters the systemic circulation*
- c. remove the drug from the body
- d. All above

426-If 100mg of the drug is administered orally and 70mg of this drug is absorbed unchanged, the bioavailability is:

- a. 100%
- b. 10%
- c. 70%*
- d. 0%

427-The nervous system is divided into:

- a. central nervous system and peripheral nervous system*
- b. peripheral nervous system only
- c. brain
- d. All of the above

428-Epinephrine is indicated for:

- a. anaphylactic shock*
- b. asthma
- c. nasal decongestant
- d. All above

429-Which of the following drug can cause addiction?

- a. barbiturate*
- b. tetracycline
- c. iodine
- d. furosemide

430- All of these drugs are used for general anesthesia except:

- a. nitrous oxide
- b. diazepam
- c. metronedazol*
- d. morphin

431- All of these drugs are used in epilepsy except:

- a. carbamazepine
- b. furosemide*
- c. diazepam
- d. phenytoin

432-Captopril:

- a. vasodilators
- b. used for treatment of congestive heart failure*
- c. a and b
- d. None of the above

433- All of these medications are anti-hypertensive except:

- a. amlodipine
- b. atenolol
- c. timolol
- d. ferrous gluconate*

434-Agents used in anemia:

- a. ferrous gluconate
- b. vitamin B12
- c. folic acid
- d. All of the above*

435-Anticoagulants:

- a. reduce the formation of fibrin clots
- b. heparin
- c. warfarin
- d. All of the above*

436-Anti platelet drugs :

- a. increase the bleeding time
- b. aspirin
- c. a and b*
- d. cloxacillin

437-Drugs used for treatment of hyperlipidemias:

- a. lovastatin*
- b. mebendazole
- c. lorazepam
- d. None of the above

438-Drugs used in treatment of asthma:

- a. aminophylline
- b. beclomethasone
- c. cimetidine
- d. a and b*

439-Penicillin and cephalosporin:

- a. are the major antibiotics that inhibit bacterial cell wall synthesis
- b. called beta lactams
- c. the microbial resistance poses constant challenge to use
- d. All of the above*

440-Amoxicillin and ampicillin:

- a. cell wall synthesis inhibitors
- b. used for treatment of infections caused by gram+ve and gram-ve
- c. contraindicated in patient with penicillin hypersensitivity
- d. All of the above*

441-Which of the following statements are correct about cephalosporin?

- a. they are classified for 4th generations
- b. do not used for treatment urinary tract infection
- c. cefoxitin is a cephalosporin derivatives
- d. a and c*

442-Protein synthesis inhibitors:

- a. tetracyclin
- b. gentamycin
- c. clarithromycin
- d. All of the above*

443-Ciprofloxacin:

- a. it is a fluoroquinolones
- b. it is bactericidal
- c. used for treatment of respiratory and urinary tract infection
- d. All above *

444-Drugs used for treatment of tuberculosis:

- a. isoniazide
- b. rifampin
- c. ethambutol
- d. All of the above*

445-Which of the following is not anti fungal agent:

- a. naproxen* NSAID
- b. amphotericin B
- c. miconazol
- d. nystatin

446-Which of the following is not anti viral agent:

- a. amantadine
- b. acyclovir
- c. ribovirin
- d. cephalixin*

447-Which of the following drugs are disinfectants:

- a. chlorohexidine
- b. cresol
- c. iodine
- d. All of the above*

448-Drugs used for treatment of allergies:

- a. chlorpheniramine maleate
- b. diphenhydramine hydrochloride
- c. indomethacin
- d. a and b*

449-Which of the following drugs are used for constipation?

- a. bisacodyl
- b. magnesium sulphate
- c. castor oil
- d. All of the above*

450-Thrombocytopenia means:

- a. Increase platelets
- b. Decrease Platelets*
- c. Decrease RBCs
- d. Increase RBCs

451-Generic name of Aldacton:

- a. aldhyde
- b. spironolactone*
- c. aldosterone
- d. phenylephrine

452-tochopherol is

- a. Vit K
- b. Vit E*
- c. B complex
- d. VitC

453-Patient received Amiloride + Captopril the expected Side effect is:

- a. hypercalcemia
- b. hyponatremia
- c. hyperkalemia*
- d. hypokalemia

454-Carviststin withdrawal from the market because it causes:

- a. rhabdomyolysis*
- b. osteomyelitis
- c. leukemia
- d. CNS.

455-Which of the following have local effect?

- a. Insulin
- b. spironolactone
- c. lidocaine
- d. Non of the above

456-sulphinpyrazone is:

- a. Analgesic
- b. Anti inflammatory
- c. Antipyretic
- d. Stimulate uric secretion*(uricosoric)

457-WFIcharacterizedby:

- a. Free from sodium
- b. Free from Ca and K
- c. Pyrogen free
- d. All above *

458-Tween 20 is:

- a. deflocoulant
- b. lipohpilic surfactant*
- c. preservative
- d. hydrophilic surfactant

459-Drug used intreatment of Hypotension:

- a. Dolobid (Diflunisal) NSAID
- b. doloxene (Dextropropoxyphene), opoid
- c. dilitiazem
- d. dopamine*

460-Normal heart beat:

- a. 100-140
- b. 60-100*
- c. 120-140
- d. 100-200

461-Crystalline Insulin is:

- a. Cloudy*
- b. Clear
- c. Unclear
- d. Nonabove

462-IncreaseTemperature means:

- a. Hypertension
- b. Hyperthermia*
- c. Hypothermia
- d. Thermal

463-Increase Heart rate:

- a. Tachycardia*
- b. bradycardia
- c. arrhythmia CHF

464-Decrease Heart rate:

- a. tachycardia
- b. bradycardia*
- c. arrhythmia CHF

465-Normal saline conc.:

- a. 9%
- b. 4.5%
- c. 0.9%*
- d. 90%

466-Intrathecal means injection into:

- a. Joints
- b. Spinal cord*
- c. Heart
- d. Muscle

467-antilice's:

- a. Topical steroid
- b. antibiotics
- c. parathion*
- d. non above

468-Most common drops:

- a. Solution*
- b. Suspension
- c. Emulsion
- d. Oint.

469-Sedating anti histaminic acting on:

- a. H1 receptor*
- b. H2 receptor
- c. H3 receptor
- d. H4 receptor

470-Noradrenalin affects all these receptors except:

- a. β_1
- b. α_1
- c. α_2
- d. β_2 *

471-Megaloplastic anemia is a Side effect of:

- a. mediazolam
- b. ketamin
- c. Nitrousoxide*
- d. Sevoflurane

472-Suitable volume for ophthalmic preparation is:

- a. 1 ml
- b. 50 microliters
- c. 10 microliters
- d. 5 ml*

473-Methadone used in:

- a. Severpain*
- b. Morphine dependence*
- c. Euphoria
- d. Respiratory depression

474-H₂O+Nacl final will be:

- a. Suspension
- b. Emulsion
- c. Solution*
- d. Lotion

475-Which of the following are Anti cholenergic?

- a Atropine
- b. Homatropine
- c. Scopolamine
- d. All above *

476-Generic name of Vasotic:

- a. Ramipril
- b. Enalapril*
- c. Captopril
- d. Lisinopril

477-Trade name of Clindamycine:

- a. Dalacin*
- b. Ceporex
- x. Kl acid
- d. Vibramycin

478-Which of the following is Catecholamine?

- a. Epinephrine
- b. Norepinephrine
- c. Serotonine
- d. Dopamine
- e. All above *

479-Generic name of Renetic:

- Captopril
- Lisinopril
- Prindopril
- Enalpril*

480-β blocker used in:

- a. treatment of Arrhythmia
- b. treatment of Hypertension
- c. treatment of Angina pectoris
- d All above *

481-Cefixime is:

- a. First generation Cephalosporin's
- b. Second generation Cephalosporin's
- c. Third generation Cephalosporin's*
- d. Fourth generation Cephalosporin's

482-Cefoxitin is:

- a. First generation Cephalosporin's
- b. Second generation Cephalosporin's*
- c. Third generation Cephalosporin's
- d. Fourth generation Cephalosporin's

483-Maximum dose of Aspirin:

- a. 300mg
- b. 5000mg
- c. 4gm*
- d. 1000mg

484-Cinnarizine used in:

- a. Hypertension
- b. Depression
- c. Peptic ulcer
- d. Prophylaxis from migraine
(Mènière's disease)

485-Generic name of Cialis:

- a. sildenafil
- b. pravastatin
- c. vardenafil
- d. tadalafil*

486-Trade name of Benazepril:

- a. Capoten
- b. Renitec
- c. Zestril
- d. Cibacen*

487-Generic name of Capril:

- a. Diclofenac potassium
- b. Captopril*
- c. Carbamazepine
- d. Cefixime

488-Generic name of Zestril

- a. Prendopril
- b. Benazepril
- c. Lisinopril*
- d. Quinapril

489-Generic name of Elyzol:

- a. Metronidazol*
- b. Metrizamide
- c. Loperamide
- d. Allopurinol

490-Trade name of Prindopril:

- a. Capozide
- b. Amoxil
- c. Ceporex
- d. Coversyl*

491-Co-Renitec is:

- a. Captopril + Hydrochlorothiazide
- b. Enalapril + Chlorothiazide
- c. Enalapril + Hydrochlorothiazide*
- d. Lisinopril + Indapamide

492-Generic name of Acuitel is:

- a. Ramipril
- b. Lisinopril
- c. Fosinopril
- d. Quinapril*

493-Chlorothiazide is:

- a. Loop diuretics
- b. Potassium sparing diuretics
- c. Osmotic diuretics
- d. Thiazide diuretics*

494-Cefaclor is:

- a. Cephalosporine Antibiotic*
- b. β blocker
- c. Aminoglycoside Antibiotic
- d. Macrolide Antibiotic
- e. cough suppressant

495-Atenolo is:

- a. Cephalosporine Antibiotic
- b. β blocker *
- c. Aminoglycoside Antibiotic
- d. Macrolide Antibiotic
- e. Cough suppressant

496-Clarithromycin is:

- a. Cephalosporine Antibiotic
- b. β blocker
- c. Aminoglycoside Antibiotic
- d. Macrolide Antibiotic*
- e. Cough suppressant

497-Amikacin is:

- a. Cephalosporine Antibiotic
- b. β blocker
- c. Aminoglycoside Antibiotic*
- d. Macrolide Antibiotic
- e. Cough suppressant

498-Pholcodine is:

- a. Cephalosporine Antibiotic
- b. β blocker
- c. Aminoglycoside Antibiotic
- d. Macrolide Antibiotic
- e. Cough suppressant*

499-Non OTC drug:

- a. Malox
- b. Bisacodyl
- c. Acetamenophen
- d. Acetyl salicylic acid
- e. Tetracycline*

500-Carbidopa used in treatment of :

- a. Parkinson's*
- b. Anxiety
- c. Leprosy
- d. Amoeba
- e. Epilepsy
- f. Muscle inflammation

501-Amiodarone used as Oral Anti coagulant

- a. Heparin Antagonist
- b. Anti plattlet
- c. Anti arrhythmic*
- d. Parenteral Anticoagulant

502-Ibuprofen used intreatment of :

- a. Parkinson's
- b. Anxiety
- c. Leprosy
- d. Amoeba
- e. Epilepsy
- f. Muscle inflammation*

503-Metronidazol used intreatment of :

- a. Parkinson's
- b. Anxiety
- c. Leprosy
- d. Amoeba*
- e. Epilepsy
- f. Muscle inflammation

504-Diazepam used intreatment of :

- a. Parkinson's
- b. Anxiety*
- c. Leprosy
- d. Amoeba
- e. Epilepsy
- f. Muscle inflammation

505-Phyntoin used intreatment of :

- a. Parkinson's
- b. Anxiety
- c. Leprosy
- d. Amoeba
- e. Epilepsy*
- f. Muscle inflammation

506-warfarine used as:

- a. Oral Anticoagulant*
- b. Heparin Antagonist
- c. Antiplattlet
- d. Anti arrhythmic
- e. Parenteral Anticoagulant

507-Protamin sulphate used as:

- a. Oral Anticoagulant
- b. Heparin Antagonist*
- c. Anti plattlet
- d. Anti arrhythmic
- e. Parenteral Anticoagulant

508-Heparin used as:

- a. Oral Anticoagulant
- b. Heparin Antagonist
- c. Anti plattlet
- d. Anti arrhythmic
- e. Parenteral Anticoagulant*

509-Aspirin used as:

- a. Oral Anticoagulant
- b. Heparin Antagonist
- c. Anti plattlet*
- d. Anti arrhythmic
- e. Parenteral Anticoagulant

510-Ca channel blocker :

- a. Enalapril
- b. Isradepine*
- c. Metoprolol
- d. Pergolide
- f. Phenyl Ephrine

511-Angiotensin converting Enzyme Inhibitor:

- a. Enalapril*
- b. Isradepine
- c. Metoprolol
- d. Pergolide
- e. Phenyl Ephrine

512- α Agonist:

- a. Enalapril
- b. Isradepine
- c. Metoprolol
- d. Pergolide
- e. Phenylephrine*

514-which type of Insulin taken orally:

- a. NPH
- b. Mixtard
- c. Actrapid
- d. Insulin not taken orally*

515-Regular Insulin characters:

- a. clear
- b. colorless
- c. watery
- d. All of the above*

516-Fosinopril is:

- a. Staril*
- b. Capoten
- c. Acutil
- d. Renetic

517-All of these Anticoagulant except:

- a. heparin
- b. warfarin
- c. ancord
- d. clopidogrel*

518-Ticlid® is:

- a. Dipyridamol
- b. Aspirin
- c. Ticlodipine*
- d. Enoxaparine

519-The meaning of drug that is absolutely contraindicated in pregnancy by FDA:

- a. Pregnancy Category A
- b. Pregnancy Category B
- c. Pregnancy Category C
- d. Pregnancy Category D
- e. Pregnancy Category X*

520-Acetab is:

- a. Ramipril
- b. Captopril*
- c. lisinopril
- d. Enalapril

521-Which of these is/are catecholamine's:

- a. Epinephrine
- b. Norepinephrine
- c. Dopamine
- d. Dobutamine
- e. Serotonin
- f. All of them*

522-Which of the following can't taken with out prescription:

- a. H2 antagonist
- b. Antibiotic
- c. Narcotic drug
- d. Corticosteroids
- e. All of them*

523- All of these are used in T.B except:

- a. I.N.H
- b. Cycloserine
- c. Rifampin
- d. ethambutol
- e. Cyclosporin*

524-Bethanecol is:

- a. Muscarinic agonist*
- b. Adrenergic antagonist
- c. Muscarinic Antagonist
- d. Adrenergic agonist

525-Which is not parasympatholytic:

- a. Hyocine
- b. Omeprazol*
- c. Atropine
- d. Benztropine
- e. None of them

526- Imigran used for:

- a. treatment of Amoeba
- b. treatment of diharrea
- c. treatment of migraine*
- d. treatment of Indigestion

(Sumatriptan is serotonineagonist, used for the acute treatment of migraine)

527-Natural emulsifying agent is:

- a. Span
- b. Acacia*
- c. Tween
- d. All above

528-Injection inspinal fluid called:

- a. Intrathecal injection*
- b. Intra muscular
- c. Intra pretonial
- d. Intra articular

529-which of these is fibrinolytic:

- a. Urikinase
- b. Streptokinase
- c. Alteplase
- d. All of them*

530-Terfenadine is:

- a. Antibiotic
- b. Antifungal
- c. Antiemetic
- d. Antihistamine*

531-Cilazapril trade name is:

- a. Enalapril
- b. Capoten
- c. Vascase*
- d. Zestril

532-Zelaic acid used for:

- a. Diarrhea
- b. Anti coagulant
- c. Antacid
- d. Acne*

533-Propantheline bromide is:

- a. Antiprotozoal
- b. Antibiotic
- c. Antimuscarinic*
- d. Antihistaminic

534-Acenocoumarol is

- a. Anticoagulant*(oral)
- b. Antiarrhythmic
- c. Antihypertensive
- d. Antiplatelets

535-Daptomycin is:

- a. Antiviral
- b. Antibiotic*
- c. Antifungal
- d. a and b

536-How many grams of water are used to prepare 60g of potassium acetate solution have concentration 5%w/w:

- a. 300gm
- b. 57gm*
- c. 3gm
- d. 5.7gm

537-How many grams of substance X are needed to prepare 100cc from solution 1:1000?

- a. 10mg
- b. 0.01mg
- c. 100mg*(0.1g)
- d. 1000mg

538-Best route of administration for Chloramphenicol is:

- a. Oral
- b. Parenteral
- c. Topical*(eye,ear,skin)
- d. S.c
- e. Inhalation

539-Most of drugs are:

- a. Weak acids
- b. Weak bases*
- c. Strong acids
- d. Weak salts*
- e. Strong salts

540-Malaria is:

- a. Bacteria
- b. Fungus
- c. Virus
- d. Protozoa*

541-One of these substances not medically used :

- a. Morphine
- b. Codeine
- c. Heroin*
- d. Phenobarbital

542-- Which of the following is not a Side effect of codeine?

- a. Miosis
- b. Nausea
- c. Diarrhea*
- d. Respiratory depression
- e. Addiction

543-Epinephrine is not given orally because:

- a. It is inactivated in the gastric mucosa*
- b. Local vasoconstriction inhibits absorption
- c. It is rapidly inactivated in circulation
- d. None of the above
- e. All of the above

544-Various opiates may be used as All of the following except :

- a. Analgesics
- b. Anti-inflammatories*
- c. Antidiarrheals
- d. Aids to anesthesia
- e. Antitussive

545-Which of the following action is not seen with sympathomimetics?

- a. Pupil constriction*
- b. Increased heart rate
- c. Bronchiole dilation
- d. Systemic blood vessel constriction

546-Which of the following diuretics do not cause potassium loss?

- a. Furosemide
- b. Chlorthalidone
- c. Triamterene*
- d. Hydrochlorothiazide
- e. Metolazole

547-Which of the following beta. adrenergic agents is not short acting?

- a. Pirbuterol
- b. Salmeterol*
- c. Terbutaline
- d. Albuterol
- e. None of the above

548-Possible treatment of Hypoglycemia does not include:

- a. Insulin *
- b. Candy
- c. Glucose
- d. Fruitjuce
- e. Glucagon

549-Which of the following drugs is not aTricyclic Anti depressant?

- a. Trimipramine
- b. Maprotiline*(tetracyclic anti depressant)
- c. Nortryptiline
- d. Imipramine HCL

550-Captopril and Enalapril do all the following except:

- a. Increase rennin concentration in blood
- b. Inhibit an enzyme
- c. Competively blocks Angiotensin|| at it'sreceptors*
- d. Raises the blood Potassium concentration

551-Which of the following factors affect the distribution of a drug?

- a. Lipidsolubility
- b. Plasma protein binding
- c. Polarity
- d. Molecularsize
- e. All of the above*

552-Salicylic acid is used primarily as an(a)

- a. Analgesic*
- b. Anti pyretic
- c. Cough suppressant
- d. Uricosuric agent
- e. Keratolytic agent*

553-Which of the following factors affect the distribution of a drug?

- a. Lipid solubility
- b. Plasma protein binding
- c. Polarity
- d. Molecular size
- e. All of the above*

554-Most drugs are

- a. Strong electrolytes
- b. Weak electrolytes*
- c. None lectrolytes
- d. Highly ionic
- e. None of the above

555-The most common disintergrator in compressed tablets is

- a. Dextrose
- b. Lactose
- c. Starch*
- d. Potassium bitartrate
- e. Powdered sucrose

556-Which of the following is found in vitamin B12?

- a. Magnesium
- b. Nickel
- c. Iron
- d. Cobalt*
- e. Manganese

557-Gums are used in tableting primarily as

- a. Disintegrators
- b. Glidants
- c. Lubricants
- d. Binding agents*
- e. Both b and c

558-Which of the following is classified as fat soluble?

- a. Vitamin d*
- b. Niacin
- c. Ascorbic acid
- d. Thiamine hydrochloride
- e. Riboflavin

559-Ophthalmic solutions should be formulated to include which of the following ?

- a. Sterility
- b. Isotonicity
- c. Buffering
- d. All above *



560-Advantages of tablets over liquid dose forms include the following

- a. Enhanced stability
- b. Ease of administration
- c. Greater accuracy of dosing
- d. All above *

561-Activated charcoal is used in some antidotes because of which of its properties?

- a. Neutralizing
- b. Emetic
- c. Absorptive
- d. Adsorptive*
- e. Stabilizing

562-parkinson is probably due to

- a. too little dopamine in the brain*
- b. too little levodopa in the brain
- c. too little acetylcholine in the brain
- d. too much levodopa in the brain
- e. too much dopamine in the brain

563-Cyclosporine is used for

- a. allergies
- b. angina
- c. prevention of transplant rejection*
- d. steroids deficiency
- e. treating lead poisoning

564-pantoprazole is used primarily to treat

- a. gastric hyperacidity*
- b. Hypertension
- c. cardiac insufficiency
- d. gout
- e. migraine headache

565-citalopramisa (an)

- a. diuretic
- b. cardiogenic
- c. anti depressant* **SSRI**
- d. anti-inflammatory
- e. an thelmintic

566-Ultra short acting barbiturates are used primarily as

- a. Sedatives
- b. Hypnotics
- c. Antispasmodic agents
- d. Anti-parkinsoni anagents
- e. Preanesthetic agents*

567-Orlistat is used as a(an)

- a. Narcotic antagonist
- b. Narcotic analgesic
- c. Weight loss agent*
- d. Antiepileptic
- e. Anesthetic

568-Lidocaine is used as a local anesthetic or as a (an)

- a. General anesthetic
- b. Antipruritic
- c. Preanesthetic
- d. Antiarrhythmic*
- e. Analgesic

569-Which of the following is classified as a cholinergic antagonist?

- a. Acetylcholine
- b. Neostigmine
- c. Atropine*
- d. Bethanecol
- e. Methacholine

570-Streptokinase is used to

- a. Dissolve blood clots*
- b. Treat digestive disorders
- c. Promote carbohydrate degradation
- d. Treat muscle injuries
- e. Replace pepsin

571-Gold compounds have been used to treat

- a. Worm infestations
- b. Ulcers
- c. Kidney failure
- d. Rheumatoid arthritis*
- e. Psoriasis

572-Tamoxifen is classified as a(an)

- a. Estrogen
- b. Antiestrogen*
- c. Androsterone
- d. Testosterone
- e. Thyroidhormone

573-Castor oil is classified as which type of laxative?

- a. Lubricating
- b. Anthraquinone
- c. Irritant*
- d. Stool softening
- e. Bulk producing

574-Albuterol is:

- a. Alpha receptor antagonist
- b. Alpha receptor agonist
- c. Beta receptor antagonist
- d. Beta receptor agonist*
- e. Alpha and beta receptor agonist

575-Which of the following is(are) dopaminergic antiparkinson agents?

- a. Bromocriptine(Parlodel)
- b. Pergolide(Permax)
- c. Selegiline(Eldepryl)
- d. All above *

576-Tomoxifen (Nolvadex) is an agent that can best be described as:

- a. Gonadotropin - releasing hormone analog
- b. Antiestrogen*
- c. Enstrogen
- d. Progestin
- e. Androgen

577-Which one of the following agents is classified pharmacologically as a carbonic anhydrase inhibitor

- a. Indapamide(Lozol)
- b. Acetazolamide(Diamox)*
- c. Chlorthalidone(Hygroton)
- d. Torsemide(Demadex)
- e. Amiloride(Midamor)

578-The thiazide diuretics decrease the excretion of

- a. Uric acid
- b. Urea*
- c. Sodium
- d. Bicarbonate
- e. Creatinine

579-The appropriate PH range for ophthalmic products is

- a. 2.0to3.0
- b. 4.0to6.0
- c. 6.0to8.0*
- d. 8.0to10.0

580- Side effects of isotretinoin (Roaccutane) include

- a. Dermatological reactions
- b. Depression
- c. Teratogenicity
- d. All of the above*

581-Sucral fate is used to treat

- a. Pepticulcers*
- b. Carcinoma
- c. Calciumdepletion
- d. Obesity

582-Methyldopais used as a/an

- a. Anti parkinsonism
- b. Sedative
- c. Anti hypertensive*
- d. Prolactin inhibitor

583-Lidocaine is used as

- a. General anaesthesia
- b. Antipruritic
- c. Antiarrythmic*
- d. Preanaesthetic

584-Simethicone is a\an

- a. Antacid
- b. Antiflatulent*
- c. Constipating agent
- d. Laxative

585-Which of the following drugs is the choice for the treatment of all forms of Schistosomiasis

- a. Praziquantel*
- b. Mebendazole
- c. Niclosamide
- d. Thiabendazole

586-Famotidine is classified as

- a. Anti depressant
- b. H1receptor antagonist
- c. H2 receptor antagonist*
- d. Beta2 agonist

587-Which of the following is a natural emulsifying agent

- a. Acacia*
- b. Lactose
- c. Polysorbate 20
- d. Polysorbate 80

588-Paracetamol

- a. Is not recommended for post-immunization in 2-3 month old
- b. Has an anti-inflammatory activity similar to aspirin
- c. Is an effective analgesic in severe renal colic
- d. May causes evere liver toxicity if takenin over dose*

589-for a 4 year old child, the maximum daily dose of paracetamol is

- a. 240mg
- b. 480mg
- c. 1g*
- d. 2g

590-All the following are considered to be beta-lactam type antibiotics, EXCEPT

- a. Ceftriaxone
- b. Pieracillin
- c. Ofloxacin*
- d. Imipenem

591-Which of the following drugs inhibits the formation of uric acid

- a. Aspirin
- b. Colchicine*
- c. Allopurinol
- d. Sodium salicylate

592-The most common Side effect related to the use of Aluminium containing antacids is

- a. Nausea, Vomiting
- b. Constipation*
- c. Diarrhoea
- d. GI bleeding

593-Salmeterol is a

- a. Abeta. agonist*
- b. Methylxanthine
- c. Mast-cell-stabilizer
- d. Muscarinic antagonist

594-a medicine expires" January 2001"it must not be used after

- a. 01 January ,2001
- b. 15 January ,2001
- c. 31 January ,2001*
- d. 31 December ,2001

595-Which of the following is not a beta. lactam antibiotic

- a. Aztreonam
- b. Imipenem
- c. Streptomycin*
- d. Ampicillin

596-The normal range of FASTING blood glucose is

- a. 55-75mg/dL *(60-100)
- b. 75-150mg/dL
- c. 150-180mg/dL after meal
- d. 90-110mg/dL

597-The dose of a drug is 0.5mg/kg What dose should be given to a 6yr.old child who weighs 20Kg approximately

- a. 0.003g
- b. 0.010g*
- c. 0.033g
- d. 0.05g

598-Which of the following is not indicated for use in children less than12 years old

- a. Domperidone (Motilium)
- b. Doxycycline (Vibramycin)*
- c. Chloramphenicol (Chloromycetin)
- d. Astemizole (Hismanal)

599-H2 receptor antagonist


- a. Omeprazole
- b. Sucralfate
- c. Famotidine*
- d. Metoclopramide

600-The principal toxic effect of Heparin is

- a. Bronchospasm
- b. Chills
- c. Hair loss
- d. Haemorrhage

Good Luck!


لاتنسونا من خالص الدعاء
جميع النسخ لمختلف التخصصات محفوظة على التلجرام عبر الرابط التالي
<https://t.me/medicaL02017>
للمشاركة في حل وارسال النماذج +967 733443846




نماذج قسم الصيدلة




النسخة الاخيرة مساعدين + تمرير
الجزء الثالث



نماذج اختبارات مزاولة
قبالة



نماذج اختبارات مزاولة
مختبرات



تمرير + مساعدين اطباء
بالحل وبدون حل